

QCD String Spectrum 2002

K. Jimmy Juge^a, Julius Kuti^{b *} and Colin Morningstar^c

^aInstitute for Theoretical Physics, University of Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland

^bUniversity of California at San Diego, La Jolla, CA 92093, USA

^cCarnegie Mellon University, Pittsburgh, PA 15213, USA

Results from a comprehensive new analysis on the excitation spectrum of the QCD string are presented. A rapid onset of string formation is observed in the spectrum on a length scale of 2 fm, with Dirichlet boundary conditions. The crossover from the short distance spectrum towards string excitations and an observed fine structure in the 1–3 fm range are related to effective string theory. The deficiencies of the Nambu-Goto bosonic string model in describing the observed spectrum are briefly discussed.

We report here our new comprehensive analysis of the rich low-lying energy spectrum of the excited gluon field between a static quark–antiquark ($q\bar{q}$) pair in the fundamental color representation of $SU(3)_c$. For references on earlier work, we refer to our pilot journal publication [1]. The analysis of each simulation will be detailed in forthcoming publications.

Three exact quantum numbers which are based on the symmetries of the problem determine the classification scheme of the gluon excitation spectrum in the presence of a static $q\bar{q}$ pair. We adopt the standard notation from the physics of diatomic molecules and use Λ to denote the magnitude of the eigenvalue of the projection $\mathbf{J}_g \cdot \hat{\mathbf{R}}$ of the total angular momentum $\mathbf{J}_g$ of the gluon field onto the molecular axis with unit vector $\hat{\mathbf{R}}$. The capital Greek letters $\Sigma, \Pi, \Delta, \Phi, \dots$ are used to indicate states with $\Lambda = 0, 1, 2, 3, \dots$, respectively. The combined operations of charge conjugation and spatial inversion about the midpoint between the quark and the antiquark is also a symmetry and its eigenvalue is denoted by η_{CP} . States with $\eta_{CP} = 1(-1)$ are denoted by the subscripts $g(u)$. There is an additional label for the Σ states; Σ states which are even (odd) under a reflection in a plane containing the molecular axis are denoted by a superscript $+$ ($-$). Hence, the low-lying lev-

els are labeled $\Sigma_g^+, \Sigma_g^-, \Sigma_u^+, \Sigma_u^-, \Pi_g, \Pi_u, \Delta_g, \Delta_u$, and so on. For convenience, we use Γ to denote these labels in general.

Restricted to the $R = 0.2 - 3$ fm range of a selected simulation, energy gaps ΔE above the ground state are compared to asymptotic string gaps for 18 excited states in Fig. 1. The quantity $\Delta E/(N\pi/R) - 1$ is plotted to show percentage deviations from the asymptotic string levels. The selected states correspond to $N = 1, 2, 3, 4$ string levels in the asymptotic limit. The energy gaps, far below the null lines of the plots and strongly split at fixed N , differ from the simple string gaps for $R < 2$ fm. The Nambu-Goto (NG) bosonic string does somewhat better. The spectrum with fixed end boundary conditions in D dimensions was first calculated in Ref. [2],

$$E_N = \sigma R \left[1 - \frac{D-2}{12\sigma R^2} \pi + \frac{2\pi N}{\sigma R^2} \right]^{\frac{1}{2}}, \quad (1)$$

where σ is the asymptotic string tension. The $D = 4$ choice has quantization problems and a tachyon singularity in the spectrum at $R \approx 1/3$ fm. The gaps E_N are plotted in Fig. 1 for all N values (the gray line corresponds to $N = 1$). The states Σ_u^-, Σ_g^- and Π_g' are in the worst disagreement for $R < 2$ fm. Above 2 fm, several of the levels break away from the NG formula and stay higher than the null level of the string gaps. In addition, there is a pronounced fine structure

*Talk presented by J. Kuti.

It remains a challenge to explain the precocious onset of the asymptotic value of the conformal charge.

String limit. The rapid rearrangement of the energy levels towards asymptotic string-like ordering around $R \approx 2$ fm is remarkable. For example, the states Σ_u^- and Σ_g^- break away from their respective short distance non-string degeneracies to approach string ordering for $R > 2$ fm separation. After the rapid transition, the level orderings and approximate degeneracies at large R match, without exception, those expected of the asymptotic string modes. However, the precise separation π/R of the energy levels is not observed in the spectrum. Some of the expected degeneracies are also significantly broken.

An explanation for this fine structure is expected to come from effective QCD string theory which is formulated in terms of the Lagrangian of the two-dimensional position vector ξ of the string defined in world sheet coordinates [5,6]. Symmetries of the effective Lagrangian require a derivative expansion of the form $\mathcal{L}_{\text{eff}} = \frac{1}{2}\alpha\partial_\mu\xi \cdot \partial_\mu\xi + \dots$, where the dots represent terms with four, or more derivatives and α is a dimensional constant. For low frequency excitations, the first term, which describes massless Goldstone modes (asymptotic string modes), dominates and higher dimensional operators are expected to generate perturbative fine structure in powers of R^{-1} .

The Nambu-Goto action when expanded,

$$\mathcal{L}_{\text{eff}} = \alpha\sqrt{1 + \partial_\mu\xi \cdot \partial_\mu\xi} = \frac{1}{2}\alpha\partial_\mu\xi \cdot \partial_\mu\xi + \dots, \quad (2)$$

generates higher dimensional operators with preset coefficients which cannot describe rigidity, torsion, and other physical properties of the confining flux, if they exist. Deviations of our spectrum from the NG prediction demonstrates that the coefficients in the effective QCD string action will differ significantly from those of the NG Lagrangian (a particularly interesting form was proposed in Ref. [8]).

This work was supported by the U.S. DOE, Grant No. DE-FG03-97ER40546, the U.S. National Science Foundation under Award PHY-0099450 and the European Community's Human Potential Programme, HPRN-CT-2000-00145.




Figure 2. Short distance degeneracies and crossover in the spectrum. The solid curves are only shown for visualization. The dashed line marks a lower bound for the onset of mixing effects with glueball states which requires careful interpretation.

REFERENCES

1. K.J. Juge, J. Kuti, and C. Morningstar, hep-lat/0207004.
2. J. F. Arvis, Phys. Lett. **127B**, 106 (1983).
3. N. Brambilla, A. Pineda, J. Soto, and A. Vairo, Nucl. Phys. **B566**, 275 (2000).
4. J. Kuti, Nucl. Phys. (Proc. Suppl.) **B73**, 72 (1999).
5. M. Lüscher, K. Symanzik, P. Weisz, Nucl. Phys. **B173**, 365 (1980).
6. M. Lüscher, Nucl. Phys. **B180**, 317 (1981).
7. M. Lüscher and P. Weisz, hep-lat/0207003.
8. J. Polchinski and A. Strominger, Phys. Rev. Lett. **67**, 1681 (1991).